

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

**(19) World Intellectual Property Organization
International Bureau**



**(43) International Publication Date
24 February 2005 (24.02.2005)**

PCT

(10) International Publication Number
WO 2005/017652 A2

(51) International Patent Classification⁷: G06F

(21) International Application Number: PCT/US2004/017754

(22) International Filing Date: 4 June 2004 (04.06.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/492,682	5 August 2003 (05.08.2003)	US
60/497,470	21 August 2003 (21.08.2003)	US
60/575,499	28 May 2004 (28.05.2004)	US

(71) Applicant (for all designated States except US): ROSETTA INPHARMATICS, LLC [US/US]; 401 Terry Avenue North, Seattle, WA 98109 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SCHADT, Eric,

E. [US/US]; 810 5th Street, Kirkland, WA 98033 (US). MONKS, Stephanle, A. [US/US]; 906 NE 122nd Street, Seattle, WA 98125 (US).


(74) Agents: ANTLER, Adriane, M. et al.; Jones Day, 222 East 41st Street, New York, NY 10017-6702 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

[Continued on next page]

(54) Title: COMPUTER SYSTEMS AND METHODS FOR INFERRING CAUSALITY FROM CELLULAR CONSTITUENT ABUNDANCE DATA



(57) Abstract: Methods, computer program products, and systems are provided for associating a cellular constituent with a trait T exhibited by a species. A cellular constituent i that has at least one abundance quantitative trait locus (eQTL) coincident with a respective clinical quantitative trait locus (cQTL) for the trait of interest T is identified. For each cQTL, a determination is made as to whether (i) the genetic variation of the eQTL and (ii) the variation of the trait of interest T across the plurality of organisms are correlated conditional on an abundance pattern of the cellular constituent i across the plurality of organisms. When the genetic variation of (i) one of the eQTL and (ii) the variation of the trait of interest T across the plurality of organisms are uncorrelated conditional on the abundance pattern of the cellular constituent i, the cellular constituent i is considered causal for, and is therefore associated with, the trait of interest T.



ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- without international search report and to be republished upon receipt of that report*